

**Oxin** is a natural seaweed suspension to improve overall plant growth and Crop quality thereby increasing yields. With high phosphorus levels it is ideal for use during transplanting, initial growth stage and when root growth has been stunted.

COMPOSITION	PER LITRE
Nitrogen (N)	35g/Lt
Potassium (K)	1.0g/Lt
Sulphur (S)	400mg/Lt
Phosphorus	40mg/Lt
Total Alginic Acid	10mg/Lt

#### RATE OF APPLICATION:

Ideally, Oxin should be applied in the early morning or late afternoon.


CROP	DOSAGE	APPLICATION
Seedlings	500ml / 100Lt water	Drench seedlings tray directly after drilling the seed to stimulate uniform germination and root formation.
Transplants	500ml / 100Lt water	Dip the roots of the seedling into the solution before transplant or alternatively drench the transplants just prior or after transplanting. Will decrease mortality & stimulate root growth.
Seed Coating	500ml / 10Kg seed	Apply directly to the seed and mix thoroughly to form a uniform coating on each seed. Will increase germination and stimulate initial root formation.
Vegetables	0.75Lt – 1.5Lt / 100Lt water	Apply as full cover spray during the first 4 weeks of establishment at 7 to 14 day intervals. Will stimulate growth and decrease mortality.
Potatoes	200ml / 100Lt water 4Lt / Ha 2Lt / Ha	Soak the seed potatoes into the solution before planting. Apply a second application at tuber initiation (approximately 80% emergence). Repeat application 10-14 days later.
Ornamentals	0.75Lt – 1.5Lt / 100Lt water	Apply as full cover spray at 7 to 14 days intervals throughout the growing season.
Avocado & Mango	3Lt / Ha	Apply two sprays: 1. Early flowering, and 2. At petal drop, before lignification of flower trusses. All sprays can be positioned with calcium (Ca) sprays.
Small Grains	2Lt / Ha	Apply at 1-2 node development.

Oxin should always be tested on its own on each crop and variety before applying to larger areas to ensure that there is no adverse reaction.

Batch No : 25/02/1604

Mfg Date: 29/02/2016

Expiry Date: 29/02/2018



**OXIN®**  
A Natural Seaweed Suspension

FOR AGRICULTURE USE ONLY

500ML

**USE DILUTED SOLUTION IMMEDIATELY AND DO NOT STORE OVERNIGHT**

#### FIRST AID MEASURES:

**Eye contact:** Gently flush eyes with plenty of water for at least 15 minutes. Consult a physician if eye irritation persists.

**Skin contact:** Wash contaminated immediately with water. Remove contaminated clothing. And wash before reuse. Consult a physician if skin becomes irritated.

**Ingestion:** Rinse mouth thoroughly and drink plenty of water. DO NOT INDUCE VOMIT. Seek medical attention if irritation or symptoms persist.

**Inhalation:** Move person to fresh air. Seek immediate medical attention if exposure significant and breathing is difficult.

#### HANDLING & STORAGE:

**Handling:** Use protective clothing such as gloves, overalls, hats, gumboots, facemask or protective goggles. Avoid inhalation, skin or oral contact. Do not eat, drink or smoke during application. Wash contaminated skin and clothes.

**Storage:** Store in a tightly, closed original container and in a dry, dark and well ventilated place. Keep out of reach of children, away from food, feed stuffs and animals.

DISPOSE OF CONTAINER PROPERLY AFTER USE

#### LIMITED WARRANTY:

This product is photosensitive. Manufacturer or seller makes no warranty, whether expressed or implied, concerning the use of this product other than that for the purposes indicated on the label. Neither manufacturer nor the seller should be liable for any injury or damage caused by this product as not all pesticides and fertilisers have been tested for compatibility. Liability cannot be accepted due to misuse, mishandling or any application not specifically described on the label. Efficacy of any mix will depend upon crop type and growth stage, pesticide concerned, climatic conditions, water volumes and various other factors

Manufactured for:  
Earthlee Ventures, UK  
posta@earthleeventures.com

